

AKIR-1 and IMA-2 promote nuclear import and assembly of meiotic sister chromatid cohesion proteins in *Caenorhabditis elegans* meiosis

Richard Bowman, Nathan Balukof, Talitha Ford, and Sarit Smolikove *

Supplementary tables

Figure	N2		<i>akir-1(gk528)</i>		<i>ima-1(tm7139)</i>		<i>akir-1(gk528), ima-1(tm7139)</i>	
	gonads	nuclei	gonads	nuclei	gonads	nuclei	gonads	nuclei
1A	5	1693	5	937	5	1115	5	545
1B	5	51	6	30	6	47	13	7
2B	3	712	3	486	3	455	3	273
2C	3	NA	3	NA	3	NA	3	NA
3B	3	631	3	442	3	514	3	286
3C	3	NA	3	NA	3	NA	3	NA
4B	3	1237	3	470	3	606	3	245
5B (Diakinesis -1)	20	NA	NA	NA	NA	NA	29	NA
6B	3	826	3	647	3	535	3	354
6C	3	NA	3	NA	3	NA	3	NA
7B	3	791	3	623	3	618	3	295
7C	3	NA	3	NA	3	NA	3	NA
8B	3	709	3	499	3	539	3	259
8C	3	NA	3	NA	3	NA	3	NA
S3B	3	660	3	748	3	456	3	359
S3C	3	NA	3	NA	3	NA	3	NA

Table S1: n values for data presented in Figures. n values indicate total number of germlines and total number of nuclei analyzed for each genotype for the indicated figures. However, >15 germlines were examined qualitatively for each genotype and staining to ensure that phenotypes are consistent between germlines and the phenotype is not variable.